

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 401/04, 413/04, 405/14, A61K 31/4178, A61P 19/00		A1	(11) International Publication Number: WO 00/61576
			(43) International Publication Date: 19 October 2000 (19.10.00)
(21) International Application Number: PCT/US00/09147 (22) International Filing Date: 6 April 2000 (06.04.00) (30) Priority Data: 60/128,687 9 April 1999 (09.04.99) US (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM CORPORATION [US/US]; One Franklin Plaza, Philadelphia, PA 19103 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BURGESS, Joelle, Lor- raine [US/US]; 1250 South Collegeville Road, Collegeville, PA 19426-0989 (US). CALLAHAN, James, Francis [US/US]; 1250 South Collegeville Road, Collegeville, PA 19426-0989 (US). (74) Agents: STEIN-FERNANDEZ, Nora et al.; SmithKline Beecham Corporation, Corporate Intellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).		(81) Designated States: AE, AL, AU, BA, BB, BG, BR, CA, CN, CZ, DZ, EE, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MA, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, TZ, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(54) Title: TRIARYLIMIDAZOLES <div style="text-align: center;"> <p>(I)</p> </div>			
(57) Abstract Compounds of formula (I) or a pharmaceutically acceptable salt thereof: wherein R ₁ , R ₂ and R ₃ are various substituent groups; and one of X ₁ and X ₂ is N or CR', and the other is NR' or CHR' wherein R' is hydrogen, OH, C ₁ -alkyl, or C ₃ -7cycloalkyl; or when one of X ₁ and X ₂ is N or CR' then the other may be S or O; and their use as pharmaceuticals.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

TRIARYLIMIDAZOLES

This invention relates to pyridyl substituted triarylimidazoles which are inhibitors of the transforming growth factor, ("TGF")- β signaling pathway, in particular, the phosphorylation of smad2 or smad3 by the type I or activin-like kinase ("ALK")-5 receptor, methods for their preparation and their use in medicine, specifically in the treatment and prevention of a disease state mediated by this pathway.

TGF- β 1 is the prototypic member of a family of cytokines including the TGF- β s, activins, inhibins, bone morphogenetic proteins and Müllerian-inhibiting substance, that signal through a family of single transmembrane serine/threonine kinase receptors. These receptors can be divided in two classes, the type I or activin like kinase (ALK) receptors and type II receptors. The ALK receptors are distinguished from the type II receptors in that the ALK receptors (a) lack the serine/threonine rich intracellular tail, (b) possess serine/threonine kinase domains that are very homologous between type I receptors, and (c) share a common sequence motif called the GS domain, consisting of a region rich in glycine and serine residues. The GS domain is at the amino terminal end of the intracellular kinase domain and is critical for activation by the type II receptor. Several studies have shown that TGF- β signaling requires both the ALK and type II receptors. Specifically, the type II receptor phosphorylates the GS domain of the type I receptor for TGF- β , ALK5, in the presence of TGF- β . The ALK5, in turn, phosphorylates the cytoplasmic proteins smad2 and smad3 at two carboxy terminal serines. Generally it is believed that in many species, the type II receptors regulate cell proliferation and the type I receptors regulate matrix production. Therefore, preferred compounds of this invention are selective in that they inhibit the type I receptor and thus matrix production, and not the type II receptor mediated proliferation.

Activation of the TGF- β 1 axis and expansion of extracellular matrix are early and persistent contributors to the development and progression of chronic renal disease and vascular disease. Border W.A., Noble N.A., *N. Engl. J. Med.*, Nov. 10, 1994; **331**(19):1286-92. Further, TGF- β 1 plays a role in the formation of fibronectin and plasminogen activator inhibitor-1, components of sclerotic deposits, through the action of smad3 phosphorylation by the TGF- β 1 receptor ALK5. Zhang Y., Feng X.H., Derynck R., *Nature*, Aug. 27, 1998; **394**(6696):909-13; Usui T., Takase M., Kaji Y., Suzuki K., Ishida K., Tsuru T., Miyata K., Kawabata M., Yamashita H., *Invest. Ophthalmol. Vis. Sci.*, Oct. 1998; **39**(11):1981-9.

Progressive fibrosis in the kidney and cardiovascular system is a major cause of suffering and death and an important contributor to the cost of health care. TGF- β 1 has been implicated in many renal fibrotic disorders. Border W.A., Noble N.A., *N. Engl. J. Med.*, Nov 10, 1994; **331**(19):1286-92. TGF- β 1 is elevated in acute and chronic glomerulonephritis, Yoshioka K., Takemura T., Murakami K., Okada M., Hino S., Miyamoto H., Maki S., *Lab. Invest.*, Feb. 1993; **68**(2):154-63, diabetic nephropathy, Yamamoto, T., Nakamura, T., Noble, N.A., Ruoslahti, E., Border, W.A., (1993) *PNAS* **90**:1814-1818, allograft rejection, HIV nephropathy and

angiotensin-induced nephropathy, Border W.A., Noble N.A., *N. Engl. J. Med.*, Nov. 10, 1994; **331**(19):1286-92. In these diseases the levels of TGF- β 1 expression coincide with the production of extracellular matrix. Three lines of evidence suggest a causal relationship between TGF- β 1 and the production of matrix. First, normal glomeruli, mesangial cells and non-renal cells can be induced to produce extracellular-matrix protein and inhibit protease activity by exogenous TGF- β 1 in vitro. Second, neutralizing anti-bodies against TGF- β 1 can prevent the accumulation of extracellular matrix in nephritic rats. Third, TGF- β 1 transgenic mice or in vivo transfection of the TGF- β 1 gene into normal rat kidneys resulted in the rapid development of glomerulosclerosis. Kopp J.B., Factor V.M., Mozes M., Nagy P., Sanderson N., Bottinger E.P., Klotman P.E., Thorgeirsson S.S., *Lab Invest*, June 1996; **74**(6):991-1003. Thus, inhibition of TGF- β 1 activity is indicated as a therapeutic intervention in chronic renal disease.


TGF- β 1 and its receptors are increased in injured blood vessels and are indicated in neointima formation following balloon angioplasty, Saltis J., Agrotis A., Bobik A., *Clin Exp Pharmacol Physiol*, Mar. 1996; **23**(3):193-200. In addition TGF- β 1 is a potent stimulator of smooth muscle cell ("SMC") migration in vitro and migration of SMC in the arterial wall is a contributing factor in the pathogenesis of atherosclerosis and restenosis. Moreover, in multivariate analysis of the endothelial cell products against total cholesterol, TGF- β receptor ALK5 correlated with total cholesterol ($P < 0.001$) Blann A.D., Wang J.M., Wilson P.B., Kumar S., *Atherosclerosis*, Feb. 1996; **120**(1-2):221-6. Furthermore, SMC derived from human atherosclerotic lesions have an increased ALK5/TGF- β type II receptor ratio. Because TGF- β 1 is over-expressed in fibroproliferative vascular lesions, receptor-variant cells would be allowed to grow in a slow, but uncontrolled fashion, while overproducing extracellular matrix components McCaffrey T.A., Consigli S., Du B., Falcone D.J., Sanborn T.A., Spokojny A.M., Bush H.L., Jr., *J Clin Invest*, Dec. 1995; **96**(6):2667-75. TGF- β 1 was immunolocalized to non-foamy macrophages in atherosclerotic lesions where active matrix synthesis occurs, suggesting that non-foamy macrophages may participate in modulating matrix gene expression in atherosclerotic remodeling via a TGF- β -dependent mechanism. Therefore, inhibiting the action of TGF- β 1 on ALK5 is also indicated in atherosclerosis and restenosis.

TGF- β is also implicated in peritoneal adhesions Saed G.M., Zhang W., Chegini N., Holmdahl L., and Diamond MP., *Wound Repair Regeneration*. **7**(6):504-510, 1999 Nov-Dec. Therefore, inhibitors of ALK5 would be beneficial in preventing peritoneal and sub-dermal fibrotic adhesions following surgical procedures.

Surprisingly, it has now been discovered that a class of 2-pyridyl substituted triarylimidazoles of formula (I), function as potent and selective non-peptide inhibitors of ALK5 kinase and therefore, have utility in the treatment and prevention of various disease states mediated by ALK5 kinase mechanisms, such as chronic renal disease, acute renal disease, wound healing, arthritis, osteoporosis, kidney disease, congestive heart failure, ulcers, ocular disorders, corneal wounds, diabetic

nephropathy, impaired neurological function, Alzheimer's disease, trophic conditions, atherosclerosis, peritoneal and sub-dermal adhesion, any disease wherein fibrosis is a major component, including, but not limited to lung fibrosis and liver fibrosis, and restenosis.

- 5 According to the invention there is provided a compound of formula (I) or a pharmaceutically acceptable salt thereof:



(I)

- 10 wherein R_1 is naphthyl, anthracenyl, or phenyl optionally substituted with one or more substituents selected from the group consisting of halo, C_{1-6} alkoxy, C_{1-6} alkylthio, C_{1-6} alkyl, $-O-(CH_2)_n-Ph$, $-S-(CH_2)_n-Ph$, cyano, phenyl, and CO_2R , wherein R is hydrogen or C_{1-6} alkyl and n is 0, 1, 2 or 3; or R_1 is phenyl fused with an aromatic or non-aromatic cyclic ring of 5-7 members wherein said cyclic ring optionally
 15 contains up to two heteroatoms, independently selected from N, O and S;
 R_2 is H, $NH(CH_2)_n-Ph$ or $NH-C_{1-6}alkyl$, wherein n is 0, 1, 2 or 3;
 R_3 is CO_2H , $CONH_2$, CN, NO_2 , $C_{1-6}alkylthio$, $-SO_2-C_{1-6}alkyl$, $C_{1-6}alkoxy$, $SONH_2$, $CONHOH$, NH_2 , CHO, CH_2OH , CH_2NH_2 , or CO_2R , wherein R is hydrogen or $C_{1-6}alkyl$;
 20 and

one of X_1 and X_2 is N or CR' , and the other is NR' or CHR' wherein R' is hydrogen, OH, $C_{1-6}alkyl$, or $C_{3-7}cycloalkyl$; or when one of X_1 and X_2 is N or CR' then the other may be S or O.

- As used herein, the double bond indicated by the dotted lines of formula (I),
 25 represent the possible tautomeric ring forms of the compounds falling within the scope of this invention. It will be understood that when one of X_1 and X_2 is carbon and the other is nitrogen, then the double bond could be either to the carbon or the nitrogen. When X_1 and X_2 are both carbon, then the double bond could be to either X_1 or X_2 . When X_1 and X_2 are both nitrogen, then the double bond is to the
 30 unsubstituted nitrogen.

- Preferably R_1 is optionally substituted naphthyl or phenyl. More preferably R_1 is phenyl optionally substituted with one or more substituents selected from the group consisting of halo, $C_{1-6}alkoxy$, $C_{1-6}alkylthio$, and phenyl; or R_1 is phenyl fused with an aromatic or non-aromatic cyclic ring of 5-7 members wherein said cyclic
 35 ring optionally contains up to two heteroatoms, independently selected from N, O and S, for example R_1 represents benzo[1,3]dioxolyl, 2,3-dihydrobenzo[1,4]dioxinyl, benzoxazolyl, benzothiazolyl, benzo[1,2,5]oxadiazolyl, benzo[1,2,5]thiadiazolyl or dihydrobenzofuranyl.

Preferably, R₂ is positioned ortho to the nitrogen of the pyridyl ring.
More preferably R₂ is hydrogen.

Preferably R₃ is CO₂H, CONH₂, CN, NO₂, SONH₂, CONHOH, NH₂, CHO, CH₂OH or CH₂NH₂.

- 5 Preferably one of X₁ and X₂ is N or CR', and the other is NR' or CHR' wherein R' is hydrogen, C₁₋₆alkyl, or C₃₋₇cycloalkyl, provided that at least one of X₁ and X₂ is N or NR'; or one of X₁ and X₂ is N and the other is O. More preferably one of X₁ and X₂ is N and the other is NR'.

Preferably each R' is hydrogen.

- 10 Specific compounds of the invention which may be mentioned include the following and pharmaceutically acceptable salts thereof:

4-[4-(4-Fluorophenyl)-5-(2-pyridyl)-1-hydroxy-1*H*-imidazol-2-yl]benzonitrile;

- 4-[4-(4-Fluorophenyl)-5-(2-pyridyl)-1*H*-imidazol-2-yl]benzonitrile;
15 4-[4-(4-Fluorophenyl)-5-(2-pyridyl)-1*H*-imidazol-2-yl]benzoic acid;
Methyl 4-[4-(4-fluorophenyl)-5-(2-pyridyl)-1*H*-imidazol-2-yl]benzoate;
Ethyl 4-[4-(4-fluorophenyl)-5-(2-pyridyl)-1*H*-imidazol-2-yl]benzoate
4-(4-Benzo[1,3]dioxol-5-yl-1-hydroxy-5-pyridin-2-yl-1*H*-imidazol-2-yl)benzonitrile;
20 4-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1*H*-imidazol-2-yl)benzonitrile;
4-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1*H*-imidazol-2-yl)benzoic acid;
2-[4-Benzo[1,3]dioxol-5-yl-2-(4-nitrophenyl)-1*H*-imidazol-5-yl]pyridine;
3-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1*H*-imidazol-2-yl)phenylamine;
4-[4-(4-Fluorophenyl)-2-(4-nitrophenyl)-1*H*-imidazol-5-yl]pyridine;
25 4-[4-(4-Fluorophenyl)-5-pyridin-2-yl-1*H*-imidazol-2-yl]phenylamine
4-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1*H*-imidazol-2-yl)phenyl]methanol;
4-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1*H*-imidazol-2-yl)benzamide;
4-[4-(2,3-Dihydro-benzo[1,4]dioxin-6-yl)-5-pyridin-2-yl-1*H*-imidazol-2-yl]-
30 benzonitrile;
4-[4-(2,3-Dihydro-benzo[1,4]dioxin-6-yl)-5-pyridin-2-yl-1*H*-imidazol-2-yl]benzamide;
4-[4-(2,3-Dihydro-benzofuran-5-yl)-5-pyridin-2-yl-1*H*-imidazol-2-yl]benzamide;
35 3-[4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1*H*-imidazol-2-yl]benzonitrile;
4-[4-(2,3-Dihydro-benzofuran-6-yl)-5-pyridin-2-yl-1*H*-imidazol-2-yl]benzonitrile;
4-[4-(2,3-Dihydro-benzofuran-6-yl)-5-pyridin-2-yl-1*H*-imidazol-2-yl]benzamide;
40 3-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1*H*-imidazol-2-yl)benzoic acid;
4-[4-(4-Methoxyphenyl)-5-(2-pyridyl)-1*H*-imidazol-2-yl]benzonitrile;
4-[4-(2,2-Difluoro-benzo[1,3]dioxol-5-yl)-5-pyridin-2-yl-1*H*-imidazol-2-yl]benzamide;

4-[4-(2,3-Dihydro-benzo[1,4]dioxin-6-yl)-1-methyl-5-pyridin-2-yl-1*H*-imidazol-2-yl]benzamide;

4-[5-(2,3-Dihydro-benzo[1,4]dioxin-6-yl)-1-methyl-4-pyridin-2-yl-1*H*-imidazol-2-yl]benzamide;

- 5 4-(5-Benzo[1,3]dioxol-5-yl-4-pyridin-2-yl-oxazol-2-yl)benzonitrile;
4-(5-Benzo[1,3]dioxol-5-yl-4-pyridin-2-yl-oxazol-2-yl)benzamide; and
4-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1*H*-pyrrol-2-yl)benzamide.

Suitable, pharmaceutically acceptable salts of the compounds of formula (I) include, but are not limited to, salts with inorganic acids such as hydrochloride, sulfate, phosphate, diphosphate, hydrobromide, and nitrate, or salts with an organic
10 acid such as malate, maleate, fumarate, tartrate, succinate, citrate, acetate, lactate, methanesulfonate, p-toluenesulfonate, palmitate, salicylate, and stearate.

Some of the compounds of this invention may be crystallised or recrystallised from solvents such as aqueous and organic solvents. In such cases solvates may be
15 formed. This invention includes within its scope stoichiometric solvates including hydrates as well as compounds containing variable amounts of water that may be produced by processes such as lyophilisation.

Certain of the compounds of formula (I) may exist in the form of optical isomers, e.g. diastereoisomers and mixtures of isomers in all ratios, e.g. racemic
20 mixtures. The invention includes all such forms, in particular the pure isomeric forms. The different isomeric forms may be separated or resolved one from the other by conventional methods, or any given isomer may be obtained by conventional synthetic methods or by stereospecific or asymmetric syntheses.

Since the compounds of formula (I) are intended for use in pharmaceutical compositions it will readily be understood that they are each preferably provided in
25 substantially pure form, for example at least 60% pure, more suitably at least 75% pure and preferably at least 85%, especially at least 98% pure (% are on a weight for weight basis). Impure preparations of the compounds may be used for preparing the more pure forms used in the pharmaceutical compositions; these less pure
30 preparations of the compounds should contain at least 1%, more suitably at least 5% and preferably at least 10% of a compound of the formula (I) or pharmaceutically acceptable derivative thereof.

The term " C_{1-6} alkyl" as used herein whether on its own or as part of a larger group e.g. C_{1-6} alkoxy, means a straight or branched chain radical of 1 to 6 carbon
35 atoms, including, but not limited to methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl and tert-butyl.

C_{1-6} haloalkyl groups may contain one or more halo atoms, a particular C_{1-6} haloalkyl group that may be mentioned in CF_3 .

The terms "halo" or "halogen" are used interchangeably herein to mean
40 radicals derived from the elements chlorine, fluorine, iodine and bromine.

The term " C_{3-7} cycloalkyl" as used herein means cyclic radicals of 3 to 7 carbons, including but not limited to cyclopropyl, cyclopentyl and cyclohexyl.

The term "aryl" as used herein means 5- to 14-membered substituted or unsubstituted aromatic ring(s) or ring systems which may include bi- or tri-cyclic systems, including, but not limited to phenyl and naphthyl.

5 The term "ALK5 inhibitor" as used herein means a compound, other than inhibitory smads, e.g. smad6 and smad7, which selectively inhibits the ALK5 receptor preferentially over p38 or type II receptors.


10 The term "ALK5 mediated disease state" as used herein means any disease state which is mediated (or modulated) by ALK5, for example a disease which is modulated by the inhibition of the phosphorylation of smad 2/3 in the TGF- β signaling pathway.

The term "ulcers" as used herein includes, but is not limited to, diabetic ulcers, chronic ulcers, gastric ulcers, and duodenal ulcers.

15 The compounds of formula (I) can be prepared by art-recognized procedures from known or commercially available starting materials. If the starting materials are unavailable from a commercial source, their synthesis is described herein, or they can be prepared by procedures known in the art.

20 Specifically, compounds of formula (I) wherein one of X_1 and X_2 is N and the other is NH or N-OH may be prepared as illustrated in Scheme 1 for compounds wherein X_1 is N and X_2 is NH or N-OH. Using the method detailed in US 3,940,486, N-methoxy-N-methylaryl amide is alkylated with the anion generated from a 2(6)-methylpyridine to give a ketone. The ketone is treated with sodium nitrite to form the oxime which is condensed with an aldehyde and NH_4OAc to give a hydroxy imidazole. The hydroxy imidazole may then be reduced with triphenylphosphite by the method described in US 5,656,644 to give the corresponding imidazole.


25

Scheme 1

Alternatively, the ketone may be oxidised to a diketone with HBr in DMSO.


- 5 This diketone can then be condensed with a suitably substituted benzaldehyde and ammonium acetate to give the imidazole according to the method outlined in WO 98/56788 and as illustrated in Scheme 2 for compounds wherein X₁ is N and X₂ is NH.

10 Scheme 2



- 15 Non-selective alkylation of the imidazole nitrogen (using one of the procedures outlined in N. J. Liverton et al; *J. Med. Chem.*, 1999, **42**, 2180-2190) with a compound of formula L-R' wherein L is a leaving group, e.g. halo, sulfonate or triflate, will yield both isomers of the compounds where one of X₁ or X₂ is NR' wherein R' is C₁₋₆ alkyl, or C₃₋₇ cycloalkyl, as illustrated in Scheme 3, the isomers can be separated by chromatographic methods.

20


Scheme 3

5

Compounds of formula (I) wherein one of X₁ and X₂ is N and the other is O may be prepared according to Scheme 4. The oximino ketone may be reduced via catalytic hydrogen to afford the amino ketone which can be further reacted with an appropriately substituted benzoyl chloride compound. Reaction of the amide

10

product with thionyl chloride affords the oxazole product.


Scheme 4

5

Compounds of formula (I) where one of X_1 and X_2 is CH or CHR' may be prepared according to Scheme 5. A suitably substituted acetophenone and a benzaldehyde are condensed under basic conditions to afford the enone aldol product. This enone is then reacted with a pyridine-2-carboxaldehyde under sodium cyanide catalysis to afford the 1,4-diketone which is condensed with ammonium acetate to afford the pyrrole.

10

Scheme 5



During the synthesis of the compounds of formula (I) labile functional groups in the intermediate compounds, e.g. hydroxy, carboxy and amino groups, may be protected. A comprehensive discussion of the ways in which various labile functional groups may be protected and methods for cleaving the resulting protected derivatives is given in for example *Protective Groups in Organic Chemistry*, T.W. Greene and P.G.M. Wuts, (Wiley-Interscience, New York, 2nd edition, 1991).

Further details for the preparation of compounds of formula (I) are found in the examples.

The compounds of formula (I) may be prepared singly or as compound libraries comprising at least 2, for example 5 to 1,000 compounds, and more preferably 10 to 100 compounds of formula (I). Libraries of compounds of formula (I) may be prepared by a combinatorial 'split and mix' approach or by multiple parallel synthesis using either solution phase or solid phase chemistry, by procedures known to those skilled in the art.

Thus according to a further aspect of the invention there is provided a compound library comprising at least 2 compounds of formula (I) or pharmaceutically acceptable salts thereof.

The invention further provides the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for the treatment of a disease mediated by the ALK5 receptor in mammals.

The invention further provides a method of treatment of a disease mediated by the ALK5 receptor in mammals, comprising administering to a mammal in need of such treatment, a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

ALK5-mediated disease states, include, but are not limited to, chronic renal disease, acute renal disease, wound healing, arthritis, osteoporosis, kidney disease, congestive heart failure, ulcers, ocular disorders, corneal wounds, diabetic nephropathy, impaired neurological function, Alzheimer's disease, trophic
5 conditions, atherosclerosis, any disease wherein fibrosis is a major component, including, but not limited to peritoneal and sub-dermal adhesion, lung fibrosis and liver fibrosis, and restenosis.

By the term "treating" is meant either prophylactic or therapeutic therapy.

The invention further provides a method of inhibiting the TGF- β signaling
10 pathway in mammals, for example, inhibiting the phosphorylation of smad2 or smad3 by the type I or activin-like kinase ALK5 receptor, which method comprises administering to a mammal in need of such treatment, a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

The invention further provides the use of a compound of formula (I) or a
15 pharmaceutically acceptable salt thereof, in the manufacture of a medicament for inhibiting the TGF- β signaling pathway in mammals.

The invention further provides a method of inhibiting matrix formation in mammals, for example, by inhibiting the phosphorylation of smad2 or smad3 by the type I or activin-like kinase ALK5 receptor, which method comprises administering
20 to a mammal in need of such treatment, a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

The invention further provides the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for inhibiting matrix formation in mammals.

25 The compounds of formula (I) and pharmaceutically acceptable salts thereof, may be administered in conventional dosage forms prepared by combining a compound of formula (I) with standard pharmaceutical carriers or diluents according to conventional procedures well known in the art. These procedures may involve mixing, granulating and compressing or dissolving the ingredients as appropriate to
30 the desired preparation.

According to a further aspect of the present invention there is provided a pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or diluent.

35 The pharmaceutical compositions of the invention may be formulated for administration by any route, and include those in a form adapted for oral, topical or parenteral administration to mammals including humans.

The compositions may be formulated for administration by any route. The compositions may be in the form of tablets, capsules, powders, granules, lozenges,
40 creams or liquid preparations, such as oral or sterile parenteral solutions or suspensions.

The topical formulations of the present invention may be presented as, for instance, ointments, creams or lotions, eye ointments and eye or ear drops, impregnated dressings and aerosols, and may contain appropriate conventional

additives such as preservatives, solvents to assist drug penetration and emollients in ointments and creams.

The formulations may also contain compatible conventional carriers, such as cream or ointment bases and ethanol or oleyl alcohol for lotions. Such carriers may
5 be present as from about 1% up to about 98% of the formulation. More usually they will form up to about 80% of the formulation.

Tablets and capsules for oral administration may be in unit dose presentation form, and may contain conventional excipients such as binding agents, for example syrup; acacia, gelatin, sorbitol, tragacanth, or polyvinylpyrrolidone; fillers, for
10 example lactose, sugar, maize-starch, calcium phosphate, sorbitol or glycine; tableting lubricants, for example magnesium stearate, talc, polyethylene glycol or silica; disintegrants, for example potato starch; or acceptable wetting agents such as sodium lauryl sulphate. The tablets may be coated according to methods well known in normal pharmaceutical practice. Oral liquid preparations may be in the form of,
15 for example, aqueous or oily suspensions, solutions, emulsions, syrups or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives, such as suspending agents, for example sorbitol, methyl cellulose, glucose syrup, gelatin, hydroxyethyl cellulose, carboxymethyl cellulose, aluminium stearate gel or
20 hydrogenated edible fats, emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example almond oil, oily esters such as glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl *p*-hydroxybenzoate or sorbic acid, and, if desired, conventional flavouring or colouring agents.

25 Suppositories will contain conventional suppository bases, e.g. cocoa-butter or other glyceride.

For parenteral administration, fluid unit dosage forms are prepared utilizing the compound and a sterile vehicle, water being preferred. The compound, depending on the vehicle and concentration used, can be either suspended or
30 dissolved in the vehicle. In preparing solutions the compound can be dissolved in water for injection and filter sterilised before filling into a suitable vial or ampoule and sealing.

Advantageously, agents such as a local anaesthetic, preservative and buffering agents can be dissolved in the vehicle. To enhance the stability, the
35 composition can be frozen after filling into the vial and the water removed under vacuum. The dry lyophilized powder is then sealed in the vial and an accompanying vial of water for injection may be supplied to reconstitute the liquid prior to use. Parenteral suspensions are prepared in substantially the same manner except that the compound is suspended in the vehicle instead of being dissolved and sterilization
40 cannot be accomplished by filtration. The compound can be sterilised by exposure to ethylene oxide before suspending in the sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound.

The compositions may contain from 0.1% by weight, preferably from 10-60% by weight, of the active material, depending on the method of administration. Where the compositions comprise dosage units, each unit will preferably contain from 50-500 mg of the active ingredient. The dosage as
5 employed for adult human treatment will preferably range from 100 to 3000 mg per day, for instance 1500 mg per day depending on the route and frequency of administration. Such a dosage corresponds to 1.5 to 50 mg/kg per day. Suitably the dosage is from 5 to 20 mg/kg per day.

It will be recognized by one of skill in the art that the optimal quantity and
10 spacing of individual dosages of a compound of formula (I) will be determined by the nature and extent of the condition being treated, the form, route and site of administration, and the particular mammal being treated, and that such optimums can be determined by conventional techniques. It will also be appreciated by one of skill in the art that the optimal course of treatment, i.e. the number of doses of the
15 compound of formula (I) given per day for a defined number of days, can be ascertained by those skilled in the art using conventional course of treatment determination tests.

No toxicological effects are indicated when a compound of formula (I) or a pharmaceutically acceptable salt thereof is administered in the above-mentioned
20 dosage range.

All publications, including, but not limited to, patents and patent applications cited in this specification, are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

25 The following examples are to be construed as merely illustrative and not a limitation on the scope of the invention in any way. In the Examples, unless indicated otherwise, mass spectra were performed using an Hitachi Perkin-Elmer RMU-6E with chemical ionization technique (CI) or a Micromass Platform II instrument with electrospray (ES) ionization technique.

30

EXAMPLES

Example 1: 4-[4-(4-Fluorophenyl)-5-(2-pyridyl)-1-hydroxy-1H-imidazol-2-yl]benzonitrile

The title compound was prepared using the method in U.S. Patent 3,940,486 to
35 prepare 2-(t-butyl)-4-(phenyl)-N-1-hydroxy-5-(4-pyridyl)imidazole except using 2-hydroxyimino-2-(2-pyridyl)-(4-fluoro)-acetophenone and 4-cyanobenzaldehyde. ESMS (M+H)⁺=357.1.

Example 2: 4-[4-(4-Fluorophenyl)-5-(2-pyridyl)-1H-imidazol-2-yl]benzonitrile

40 The title compound was prepared using the procedure in U.S. Patent 5,656,644 (Example 1) to prepare 2-(4-cyanophenyl)-4-(4-fluorophenyl)-5-(4-pyridyl)-1H-imidazole except using 2-(4-cyanophenyl)-4-(4-fluorophenyl)-N-1-hydroxy-5-(2-pyridyl)-imidazole. ESMS (M+H)⁺=341.2.

Example 3: 4-[4-(4-Fluorophenyl)-5-(2-pyridyl)-1H-imidazol-2-yl]benzoic acid

The title compound was prepared using the procedure in U.S. Patent 5,656,644 (Example 9) to prepare 4-[4-(4-fluorophenyl)-5-(4-pyridyl)-1H-imidazol-2-yl]benzoic acid except using 4-[4-(4-fluorophenyl)-5-(2-pyridyl)-1H-imidazol-2-yl]benzonitrile. ESMS (M+H)⁺=360.1.

Example 4: Methyl 4-[4-(4-fluorophenyl)-5-(2-pyridyl)-imidazol-2-yl]benzoate

Material from Example 3 was heated at reflux in a solution of methanol/sulfuric acid for 18 h. The solution was concentrated to remove the methanol and the resulting aqueous solution was extracted with chloroform to yield the title compound. ESMS (M+H)⁺=374.2.

Example 5: Ethyl 4-[4-(4-fluorophenyl)-5-(2-pyridyl)-imidazol-2-yl]benzoate
Material from Example 3 was heated at reflux in a solution of ethanol/sulfuric acid for 18 h. The solution was concentrated to removed the ethanol and the resulting aqueous solution was extracted with chloroform to yield the title compound. ESMS (M+H)⁺=388.2.

Example 6: 4-(4-Benzo[1,3]dioxol-5-yl-1-hydroxy-5-pyridin-2-yl-1H-imidzol-2-yl)-benzonitrile

The title compound was prepared using the procedure of Example 1 except using 1-benzo[1,3]dioxol-5-yl-2-pyridin-2-ylethane-1,2-dione oxime and 4-cyanobenzaldehyde. ESMS (M+H)⁺=383.1.

Example 7: 4-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1H-imidazol-2-yl)benzonitrile

The title compound was prepared using the procedure of Example 2 except replacing 4-[4-(4-fluorophenyl)-1-hydroxy-5-pyridin-2-yl-1H-imidazol-2-yl]-benzonitrile with 4-(4-benzo[1,3]dioxol-5-yl-1-hydroxy-5-pyridin-2-yl-1H-imidazol-2-yl)-benzonitrile. ESMS (M+H)⁺=367.1.

Example 8: 4-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1H-imidazol-2-yl)benzoic acid

The title compound was prepared using the procedure of Example 3 except replacing 4-[4-(4-fluorophenyl)-5-pyridin-2-yl-1H-imidazol-2-yl]benzonitrile with 4-(4-benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1H-imidazol-2-yl) benzonitrile. ESMS (M+H)⁺=386.1.

Example 9: 2-[4-Benzo[1,3]dioxol-5-yl-2-(4-nitrophenyl)-1H-imidazol-5-yl]pyridine

The title compound was prepared using the procedure of Example 1 except using 1-benzo[1,3]dioxol-5-yl-2-pyridin-2-yl-ethane-1,2-dione oxime and 4-

nitrobenzaldehyde. ¹H NMR (250 MHz; CD₃OD) δ: 5.71 (2H, s), 6.58 (1H, d), 6.73 (2H, m), 7.03 (1H, m), 7.20-7.35 (1H, m), 7.47 (1H, m), 8.00 (4H, m), 8.31 (1H, m); m/z (ESMS): 372 [M+H].

5 **Example 10: 3-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1H-imidazol-2-yl)phenylamine**

The title compound was prepared by tin(II)chloride reduction of 2-[4-benzo[1,3]dioxol-5-yl-2-(4-nitrophenyl)-1H-imidazol-5-yl]pyridine (Example 9). m/z (ESMS): 342. [M+H].

10

Example 11: 4-[4-(4-Fluorophenyl)-2-(4-nitrophenyl)-1H-imidazol-5-yl]pyridine

The title compound was prepared using the procedure of Example 1 except using 1-(4-fluorophenyl)-2-pyridin-2-yl-ethane-1,2-dione oxime and 4-nitrobenzaldehyde.

15 m/z (ESMS): 372 [M+H].

Example 12: 4-[4-(4-Fluorophenyl)-5-pyridin-2-yl-1H-imidazol-2-yl]phenylamine

The title compound was prepared by tin(II)chloride reduction of 4-[4-(4-fluorophenyl)-2-(4-nitrophenyl)-1H-imidazol-5-yl]pyridine (Example 11). (ESMS): 342. [M+H].

20

Example 13: 4-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1H-imidazol-2-yl)phenyl]methanol

25 Methyl 4-(4-benzo[1,3]-dioxol-5-yl-4-pyridin-2-yl)benzoate (204 mg; 0.5 mmol) was dissolved in dry tetrahydrofuran (50 ml) under argon and treated dropwise with a solution of lithium hydride in tetrahydrofuran (1M; 1 ml; 1 mmol). After 1h, more lithium aluminium hydride solution (1 ml) was added and the resulting mixture stirred until tlc examination showed no starting material remaining. Saturated aq.
30 ammonium chloride solution was added, followed by ethyl acetate and the phases were separated. The organic phase was washed with water (x2), saturated brine, dried (MgSO₄) and evaporated to dryness under reduced pressure. Trituration of the residue under 60 - 80° petroleum ether gave a white solid, which was collected by filtration, washed with 60 - 80° petroleum ether and dried under reduced pressure
35 (103 mg; 54%). ¹H NMR (250 MHz, CDCl₃) δ: 2.10 - 2.80 (1H, br, exchangeable), 4.73 (2H, s), 6.02 (2H, s), 6.94 (1H, d, J = 8 Hz), 7.08 - 7.18 (3H, m), 7.41 (2H, d, J = 8 Hz), 7.53 (2H, brm), 7.92 (2H, d, J = 8 Hz), 8.53 (1H, d, J = 5 Hz); (ESMS) (M+H)⁺=372.4.

40 **Example 14: 4-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1H-imidazol-2-yl)benzamide**

4-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1H-imidazol-2-yl)benzoyl chloride hydrochloride (100 mg; 0.23 mmol) was suspended in tetrahydrofuran (5 ml) and treated with 0.88 aq. ammonia solution (1 ml) and the resulting mixture stirred at

- ambient temperature. The title compound (42 mg; 48%) was obtained as a brown powder and collected by filtration, washed with water and dried at 40°C under reduced pressure. ¹H NMR (250 MHz, CDCl₃) δ: 5.50 - 6.40 (2H, br, exchangeable), 6.03 (2H, s), 6.90 (1H, d, J = 8 Hz), 7.15 (3H, m), 7.55 (2H, brm), 7.89 (2H, d, J = 8 Hz), 8.06 (2H, d, J = 8 Hz), 8.54 (1H, brd); (ESMS) (M+H)⁺=385.4.

Example 15: 4-[4-(2,3-Dihydro-benzo[1,4]dioxin-6-yl)-5-pyridin-2-yl-1H-imidazol-2-yl]-benzonitrile

- Prepared according to the method of Example 2 from 4-[4-(2,3-dihydro-benzo[1,4]dioxin-6-yl)-1-hydroxy-5-pyridin-2-yl-1H-imidazol-2-yl]-benzonitrile. ¹H NMR (250 MHz, CDCl₃) δ: 4.32 (4H, brs), 6.94 (1H, d, J = 8.2 Hz), 7.03-7.26 (4H, m), 7.59 (2H, br. s), 7.72 (2H, d, J = 8.3 Hz), 8.07 (2H, d, J = 8.3 Hz), 8.53 (1H, br. d J= 4.6 Hz); m/z (AP⁺): 381.1 (MH⁺).

Example 16: 4-[4-(2,3-Dihydro-benzo[1,4]dioxin-6-yl)-5-pyridin-2-yl-1H-imidazol-2-yl]-benzamide

- 4-[4-(2,3-Dihydro-benzo[1,4]dioxin-6-yl)-5-pyridin-2-yl-1H-imidazol-2-yl]benzonitrile (1g) was dissolved in methanol (200 ml). Potassium carbonate was added (1g) and the resulting suspension treated dropwise with an aqueous hydrogen peroxide solution (10 ml of 30% w/v solution) at 0°C. The suspension was stirred at room temperature for 18 hours then cautiously poured into saturated sodium hydrogen sulfite solution (200 ml) at 0°C. The mixture was tested for the presence of oxidant (starch-iodide test strips) before organic solvent removal by rotary evaporation under reduced pressure. The crude product was digested in ethyl acetate (200 ml). The ethyl acetate layer was dried (MgSO₄) and evaporated to dryness under reduced pressure. The title compound was isolated by silica gel column chromatography (using a 1:9:190 ammonia:methanol: dichloromethane solution as eluent) as a beige solid (153 mg, 15%). ¹H NMR (250 MHz, CDCl₃) δ: 4.26 (4H, brs), 6.05 (1H, brs), 6.60 (1H, brs), 6.87 (1 H, brd, J = 7.7 Hz) 7.0-7.25 (3H, m), 7.53 (2H, d, J = 3.7 Hz), 7.76 (2H, d, J = 8.2 Hz), 7.95 (2H, d, J = 8.2 Hz), 8.47 (1H, d, J = 4.6 Hz); m/z (ESMS): 399 (MH⁺).

Example 17: 4-[4-(2,3-Dihydro-benzofuran-6-yl)-5-pyridin-2-yl-1H-imidazol-2-yl]-benzamide

- Prepared according to the method of Example 14 from 4-[4-(2,3-dihydro-benzofuran-6-yl)-5-pyridin-2-yl-1H-imidazol-2-yl]-benzonitrile which was prepared from 4-[4-(2,3-dihydro-benzofuran-6-yl)-1-hydroxy-5-pyridin-2-yl-1H-imidazol-2-yl]-benzonitrile according to the method of Example 2. ¹H NMR (250 MHz, CDCl₃) δ: 3.27 (2H, brt, J = 8.5 Hz), 4.63 (2H, brt, J = 8.5 Hz), 5.50-5.90 (1H, brs), 6.05-6.35 (1H, brs), 7.08-7.20 (3H, m), 7.18-7.30 (3H, m) 7.56 (2H, brs), 8.86 (2H, d, J = 8.3 Hz), 8.04 (2H, d, J = 8.3 Hz), 8.54 (1H, brd, J=4.5 Hz); m/z (ESMS): 383.5 (MH⁺).

Example 18: 3-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1*H*-imidazol-2-yl)benzonitrile

Prepared according to the method of Example 2 from 3-(4-benzo[1,3]dioxol-5-yl-1-hydroxy-5-pyridin-2-yl-1*H*-imidazol-2-yl)-benzonitrile. ¹H NMR (250MHz, CDCl₃) δ: 6.03 (s, 2H), 6.90 (d, 1H, J = 8.37 Hz), 7.15 (m, 3H), 7.57 (m, 4H), 8.16 (d, 1H, J = 7.90 Hz), 8.23 (s, 1H), 8.51 (d, 1H); m/z (AP+): 367 (M+H)⁺, 368 (M+2H)⁺.

Example 19: 4-[4-(2,3-Dihydro-benzofuran-6-yl)-5-pyridin-2-yl-1*H*-imidazol-2-yl]benzonitrile

The title compound was prepared using the procedure of Example 1 except using 1-(2,3-Dihydro-benzofuran-6-yl)-2-pyridin-2-yl-ethane-1,2-dione oxime and 4-cyanobenzaldehyde. m/z (ESMS): 365 [M+H].

Example 20: 4-[4-(2,3-Dihydro-benzofuran-6-yl)-5-pyridin-2-yl-1*H*-imidazol-2-yl]benzamide

Prepared according to the method of Example 16 from 4-[4-(2,3-dihydro-benzofuran-6-yl)-5-pyridin-2-yl-1*H*-imidazol-2-yl]benzonitrile which was prepared from 4-[4-(2,3-dihydro-benzofuran-6-yl)-1-hydroxy-5-pyridin-2-yl-1*H*-imidazol-2-yl]benzonitrile according to the method of Example 2. ¹H NMR (250 MHz, CDCl₃) δ: 3.27 (2H, brt, J = 8.5 Hz), 4.63 (2H, brt, J = 8.5 Hz), 5.50-5.90 (1H, brs), 6.05-6.35 (1H, brs), 7.08-7.20 (3H, m), 7.18-7.30 (3H, m), 7.56 (2H, brs), 8.86 (2H, d, J = 8.3 Hz), 8.04 (2H, d, J = 8.3 Hz), 8.54 (1H, brd, J=4.5 Hz); m/z (ESMS): 383.5 (MH⁺).

Example 21: 3-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1*H*-imidazol-2-yl)benzoic acid

3-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1*H*-imidazol-2-yl)-benzonitrile (600mg, 1.64mmol) was dissolved in ethanol (20ml) 2M aq. NaOH (50ml) was added and the mixture heated at reflux overnight. On cooling, the solution was acidified to pH1 with 5M aq. HCl and the precipitate collected by filtration, washed with water then diethyl ether and dried to afford a yellow/green solid. ¹H NMR (250 MHz, DMSO) δ 6.06 (s, 2H), 6.97 (d, 1H), 7.20 (dd, 3H), 7.57 (t, 1H), 7.80 (t, 1H), 7.90 (d, 1H), 8.30 (s, 2H), 8.74 (s, 1H), m/z (AP+): 385 (M+H)⁺ 386 (M+2H)⁺.

Example 22: 4-[4-(4-Methoxyphenyl)-5-(2-pyridyl)-1*H*-imidazol-2-yl]benzonitrile

The title compound was prepared using the procedure of Example 1 except using 1-(4-methoxyphenyl)-2-pyridin-2-yl-ethane-1,2-dione oxime and 4-cyanobenzaldehyde. m/z (ESMS): 353 [M+H].

Example 23: 4-[4-(2,2-Difluoro-benzo[1,3]dioxol-5-yl)-5-pyridin-2-yl-1*H*-imidazol-2-yl]benzamide

Prepared according to the method of Example 16 except using 4-[4-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-5-pyridin-2-yl-1H-imidazol-2-yl]-benzonitrile which was prepared according to the method of Example 2. m/z (ESMS): 421 [M+H].

5

Example 24: 4-[4-(2,3-Dihydro-benzo[1,4]dioxin-6-yl)-1-methyl-5-pyridin-2-yl-1H-imidazol-2-yl]benzamide

The title compound was prepared using the procedure outlined in N.J. Liverton *et al*, *J. Med Chem.* 1999, **42**, 2180-2190 via alkylation of Example 15 followed by nitrile hydrolysis as described in Example 16. ¹H NMR (250MHz, CDCl₃) δ: 3.74 (3H, s), 4.24 (4H, s), 5.30 (1H, brs), 6.25 (1H, brs), 6.74 (2H, d, J = 8Hz), 6.95 (1H, dd, J = 1,8Hz), 7.06 (1H, d, J = 1Hz), 7.26 (1H, m), 7.38 (1H, d, J = 8Hz), 7.66 (1H, m), 7.83, (2H, d, J = 7Hz), 7.91 (2H, d, J = 7Hz), 8.77 (1H, d, J = 7Hz); m/z (AP+): 413 (M+H)⁺.

15

Example 25: 4-[5-(2,3-Dihydro-benzo[1,4]dioxin-6-yl)-1-methyl-4-pyridin-2-yl-1H-imidazol-2-yl]benzamide

The title compound was prepared using the procedure outlined in N.J. Liverton *et al*, *J. Med Chem.* 1999, **42**, 2180-2190 via alkylation of Example 15 followed by nitrile hydrolysis as described in Example 16. ¹H NMR (250MHz, DMSO) δ: 3.52 (3H, s), 4.29 (4H, s), 6.92 (2H, s), 6.96 (1H, s), 7.14 (1H, m), 7.48 (1H, s), 7.81 (2H, m), 7.83 (2H, d, J = 7Hz), 8.02 (2H, d, J = 7Hz), 8.10 (1H, s), 8.34 (1H, d, J = 2Hz); m/z (AP+): 413 (M+H)⁺.

20

Example 26: 4-(5-Benzo[1,3]dioxol-5-yl-4-pyridin-2-yl-oxazol-2-yl)benzonitrile
N-(2-Benzo[1,3]dioxol-5-yl-2-oxo-1-pyridin-2-yl-ethyl)-4-cyano-benzamide (0.1g, 0.26 mmol) was dissolved in neat thionyl chloride (10 ml) and heated at reflux temperature for 2 hours. The cooled solution was cautiously added dropwise to a concentrated sodium hydroxide-ice mixture. The crude product was extracted into dichloromethane and purified using hexane-ethylacetate (1:1) as eluent to give the title compound (9 mg, 9%). ¹H NMR (CDCl₃) δ: 6.06 (2H, s), 6.84-6.99 (2H, m), 7.27-7.31 (1H, m), 7.86 (2H, d, J = 7Hz), 8.00 (2H, d, J = 7Hz), 8.04 (1H, s), 8.23 (1H, dd, J = 7 and 1 Hz), 8.42 (1H, d, J = 7Hz), 8.65 (1H, dd, J = 7 and 1 Hz); m/z (API) 368 (M+H⁺).

30

35

Example 27: 4-(5-Benzo[1,3]dioxol-5-yl-4-pyridin-2-yl-oxazol-2-yl)benzamide

Prepared from 4-(5-benzo[1,3]dioxol-5-yl-4-pyridin-2-yl-oxazol-2-yl)-benzonitrile using hydrogen peroxide and methanolic potassium carbonate according to Example 16 and isolated in 44% yield. ¹H NMR (DMSO) δ: 6.15 (2H, s), 7.08-7.16 (2H, m), 7.45-7.54 (2H, m), 7.95-7.99 (1H, m), 8.02 (1H, br. s), 8.08 (1H, br. s), 8.12-8.15 (2H, m), 8.22 (1H, dd, J = 7 and 1 Hz), 8.48 (1H, d, J = 8Hz), 8.74 (1H, d, J = 8Hz); m/z (API) 386 (M+H⁺).

40

Example 28: 4-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1H-pyrrol-2-yl)benzamide

4-(3-Benzo[1,3]dioxol-5-yl-4-oxo-4-pyridin-2-ylbutanoyl)benzamide (1.69g; 4.2mmol) and ammonium acetate (2.1g; 27.3mmol) were reacted together according to the method described in *Bioorg. Med. Chem. Letters*, 1998, 8, 2689-2694. The title compound was isolated as a brown powder from methanol (126mg; 8%).
¹H NMR (DMSO) δ: 6.04 (2H, s), 6.82 – 6.93 (4H, m), 7.18 – 7.32 (3H, m), 7.63 – 7.74 (1H, m), 7.86 – 7.96 (5H, m), 8.61 (1H, d, J = 4Hz), 11.63 (1H, brs); m/z (LCMS): 383.9 (M+H)⁺.

INTERMEDIATES

Preparation 1: 1-(2,3-Dihydro-benzo[1,4]dioxin-6-yl)-2-pyridin-2-yl-ethanol

Diisopropylamine (18.8 ml, 1.1 equiv) was dissolved in dry THF (200 ml) under argon and cooled to -78°C. n-Butyl lithium in hexane (2.5M → 53.6 ml, 1.1 equiv.) was added, followed by a dropwise addition of 2-methyl pyridine (12.0 ml, 1 equiv). The reaction mixture was stirred for 15 minutes at -78°C and was then treated with 1,4-benzodioxan-6-carboxaldehyde (20g, 121.8 mmol) in dry THF (50 ml). The reaction mixture was stirred at -78°C for 30 min then allowed to return to room temperature overnight.

Excess saturated aqueous ammonium chloride was added to the mixture followed by 200 ml EtOAc. The organic phase was separated and washed with 2M NaOH (2 x 150 ml), H₂O (3 x 100 ml), brine (100 ml) dried (MgSO₄) and evaporated to dryness. (31.6g, quantitative). ¹H NMR (400 MHz, CDCl₃) δ: 3.07 (d, 2H), 4.20 (s, 4H), 5.03 (m, 1H), 5.57 (brs, 1H), 6.81 (d, 1H), 6.85 (d, 1H), 6.94 (s, 1H), 7.09 – 7.14 (m, 2H), 7.57 (t, 1H), 8.46 (s, 1H). m/z (ES⁺): 258 (M+H)⁺, 240 (M-OH)⁺.

Preparation 2: 4-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1H-imidazol-2-yl)benzoyl chloride hydrochloride

4-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1H-imidazol-2-yl)benzoic acid (379 mg; 0.98 mmol) was suspended in dry dichloromethane (20 ml) under argon and treated sequentially with oxalyl chloride (0.1 ml) then N,N-dimethylformamide (1 drop). The mixture was stirred at ambient temperature for 1h. More oxalyl chloride (2 x 0.1 ml) was added at 1h and 2h and the mixture stirred at ambient temperature for 5h. Volatiles were removed under reduced pressure to give the title compound as an orange powder: m/z (methyl ester; API): 400.3 (M+H)⁺.

Preparation 3: 2,3-Dihydro-benzofuran-6-ol

6-Hydroxy-2H-benzofuran-3-one (3g) was suspended in anhydrous tetrahydrofuran under an argon atmosphere and cooled to 0°C. Lithium aluminium hydride (20 ml of a 1M solution in tetrahydrofuran) was added dropwise over 10 min and the reaction allowed to reach room temperature over 2 hours. The reaction was cooled to 0°C and treated dropwise with saturated ammonium chloride solution. Ethyl acetate (200 ml) was added and the mixture filtered through Celite. The ethyl

acetate layer was separated, dried (MgSO_4) and evaporated to dryness under reduced pressure. The resulting mixture of 2,3-dihydro-benzofuran-3,6-diol and benzofuran-6-ol (approximately 1:1 by 250 MHz ^1H NMR) was dissolved in a mixture of hydrochloric acid (200ml, 5M aqueous solution) and methanol (300 ml) and palladium hydroxide (0.5 g) added. The mixture was stirred under a hydrogen atmosphere for 3 hours then filtered through Celite. The organics were removed by evaporation under reduced pressure and the resulting solution neutralised with concentrated ammonia solution. The product was extracted into dichloromethane. The dichloromethane solution was dried (MgSO_4) and evaporated to dryness under reduced pressure to yield the title compound (2.5 g, 92%). ^1H NMR (250 MHz, CDCl_3) δ : 3.11 (2H, t, $J = 8.4$ Hz), 4.57 (2H, t, $J = 8.4$ Hz), 6.27-6.34 (2H, m), 6.92-7.02 (1H, m); m/z (API $^+$): 139.1($\text{M}+3\text{H}^+$).

Preparation 4: Trifluoro-methanesulfonic acid 2,3-dihydro-benzofuran-6-yl ester

2,3-Dihydro-benzofuran-6-ol (2.5 g) was dissolved in anhydrous dichloromethane under an argon atmosphere. Triethylamine (3.5 ml) was added and the reaction cooled to 0°C . A solution of trifluoromethane sulfonic anhydride (3.1 ml) in dichloromethane (10 ml) was added drop-wise over 1 minute. After 2 hours the reaction was quenched by the addition of a saturated sodium bicarbonate solution (50 ml). The reaction mixture was partitioned between water (100 ml) and dichloromethane (200 ml). The dichloromethane layer was separated, dried (MgSO_4) and evaporated to dryness under reduced pressure. The resulting residue was isolated in quantitative yield by silica gel column chromatography (using 1:3 ethyl acetate:hexane as eluent). ^1H NMR (250 MHz, CDCl_3) δ : 3.22 (2H, t, $J = 8.4$ Hz), 4.66 (2H, t, $J = 8.4$ Hz), 6.68 (1H, d, $J = 1$ Hz), 6.74 (1H, dd, $J = 8$ and 1 Hz), 7.20 (1H, d, $J = 8$ Hz); m/z (API $^+$): 271($\text{M}+3\text{H}^+$).

Preparation 5: 2,3-Dihydro-benzofuran-6-carboxylic acid

Palladium bis(triphenylphosphine) dibromide (3 g) was added to a degassed methanolic solution (30 ml) of trifluoro-methanesulfonic acid 2,3-dihydro-benzofuran-6-yl ester (5 g) and tributylamine (5 ml). The mixture was charged with carbon monoxide at 40 psi in a Berghoff pressure vessel then heated at 100°C for 72 hours. The resulting methanolic solution of 2,3-dihydro-benzofuran-6-carboxylic acid methyl ester was treated with sodium hydroxide solution (50 ml of 2M aqueous solution) and stirred at room temperature for 18 hours. The reaction mixture was acidified to pH 1 with hydrochloric acid (5M aqueous solution) and extracted with ethyl acetate (300 ml). The ethyl acetate layer was separated and the product extracted into saturated sodium bicarbonate solution (200 ml). The sodium bicarbonate solution was acidified to pH 1 with hydrochloric acid (5M aqueous solution) and the resulting precipitate dissolved in ethyl acetate. The ethyl acetate layer was separated, dried (MgSO_4) and evaporated to dryness under reduced pressure to yield the title compound (0.6g, 20%). ^1H NMR (250 MHz, CD_3OD) δ :

3.20 (2H, t, J = 8.6 Hz), 4.54 (2H, t, J = 8.6 Hz), 7.23 (1H, d, J = 8 Hz), 7.25 (1H, brs), 7.49 (1H, dd, J = 8 and 1 Hz).

Preparation 6: 2-Amino-1-benzo[1,3]dioxol-5-yl-2-pyridin-2-yl-ethanone dihydrochloride

1-Benzo[1,3]dioxol-5-yl-2-pyridin-2-yl-ethane-1,2-dione 2-oxime (2g, 7.75 mmol) was dissolved in ethanol (100 ml) and hydrochloric acid (2 ml of a 36% solution in water) was added. Palladium on charcoal catalyst (0.5 g) was added and the reaction mixture was stirred under a positive pressure of hydrogen for 1.5 hours. The catalyst was removed by filtration through Kieselguhr and the resulting filtrate evaporated to dryness under reduced pressure. Trituration under anhydrous diethyl ether gave the title compound as a pale yellow solid (1.5g, 59%); m/z (API) 257 (M+H⁺, 100%).

Preparation 7: N-(2-Benzo[1,3]dioxol-5-yl-2-oxo-1-pyridin-2-yl-ethyl)-4-cyano-benzamide

2-Amino-1-benzo[1,3]dioxol-5-yl-2-pyridin-2-yl-ethanone dihydrochloride (1g, 3 mmol) was added to a stirring solution of 4-cyano-benzoyl chloride (0.459 g, 3 mmol) in dichloromethane (10 ml). Triethylamine (2 ml, 14 mmol) was added dropwise and the reaction was stirred at ambient temperature for 18 hours. Water was (50 ml) added the organics were extracted into dichloromethane. Chromatographic purification using a dichloromethane solution of ammonia in methanol (0.5% c. ammonia : 4.5% methanol : 95% dichloromethane) yielded the title compound as a pale yellow oil (0.9g, 76%); m/z (API) 386 (M+H⁺, 100%).

Preparation 8: 4-((E/Z)-3-Benzo[1,3]dioxol-5-ylallanoyl)benzamide.

4-Acetylbenzamide (1.3g; 8mmol) was dissolved in methanol (50ml) and piperonal (1.19g; 8mmol) added, followed by 2M aqueous sodium hydroxide solution (6ml; 12mmol). The resulting mixture was left to stand at ambient temperature for 72h. The crystalline precipitate was collected by filtration, washed well with water and dried at 40°C under reduced pressure (1.51g; 64%); ν_{\max} (CH₂Cl₂) 1682, 1651, 1424 and 1265 cm⁻¹; ¹H NMR (DMSO) δ : 6.08 (2H, s), 6.96 (1H, d, J = 8Hz), 7.30 and 7.33 (1H, dd, J = 8 and 1 Hz), 7.54 (1H, br, exchangeable), 7.65 (1H, d, J = 1Hz), 7.67 (1H, d, J = 15Hz), 7.82 (1H, d, J = 15Hz), 7.98 (2H, d, J = 8Hz), 8.15 – 8.18 (3H, brd); m/z (API): 296 (M+H)⁺.

Preparation 9: 4-(3-Benzo[1,3]dioxol-5-yl-4-oxo-4-pyridin-2-ylbutanoyl)benzamide.

4-((E/Z)-3-benzo[1,3]dioxol-5-ylallanoyl)benzamide (1.47g; 4.98mmol), pyridine-2-carboxaldehyde (533mg; 4.98mmol) and sodium cyanide (24mg; 10mol%) were mixed together in N,N-dimethylformamide (30ml) and the resulting mixture heated at reflux for 18h. The reaction mixture was cooled to room temperature and poured into water with vigorous stirring then extracted with ethyl acetate. The combined organic extracts were washed with water, saturated brine, dried (MgSO₄) and

evaporated to dryness under reduced pressure to give the title compound as a black foam (1.69g; 84%); m/z (API); 403.1 (M+H)⁺.

Biological Data

5 The biological activity of the compounds of the invention may be assessed using the following assays:

Method for evaluating ALK5 kinase phosphorylation of smad3

Basic Flash-Plates (NEN Life Sciences) were coated by pipetting 100 micro liter of 0.1 molar sodium bicarbonate (pH 7.6), containing 150
10 nanograms of the fusion protein glutathion-S-transferase-smad3/100 micro liter of coating buffer. Plates were covered and incubated at room temperature for 10-24 hours. Then the plates were washed 2 times with 200 micro liter of coating buffer (0.1 molar sodium bicarbonate) and allowed to air dry for 2-4 hours.

15 For the phosphorylation reaction each well received 90 microliter containing 50 millimolar HEPES buffer (pH 7.4); 5 millimolar MgCl₂; 1 millimolar CaCl₂; 1 millimolar dithiothreitol; 100 micromolar guanosine triphosphate; 0.5 micro Ci/well gamma³³P-adenosine triphosphate (NEN Life Sciences) and 400 nanograms of a fusion protein of glutathion -S-transferase at the N-terminal end of the kinase domain of ALK5 (GST-
20 ALK5). Background counts were measured by not adding any GST-ALK5. Inhibitors of ALK5 were evaluated by determining the activity of the enzyme in the presence of various compounds. Plates were incubated for 3 hours at 30°C. After incubation the assay buffer was removed by aspiration and the
25 wells were washed 3 times with 200 microliter cold 10 millimolar sodium pyrophosphate in phosphate buffered saline. The last wash was aspirated and blotted plate dry. Plate was then counted on a Packard TopCount.

Inhibition of Matrix Markers: Western Blot Protocol

30 Data confirming activity in the enzyme assay was obtained as follows.

Cells were grown to near confluence in flasks, starved overnight and treated with TGF-beta and compounds. Cells were washed at 24 or 48 hours after treatment with ice cold phosphate buffered saline, then 500 microliter of
35 2X loading buffer was added to plate and cells were scraped and collected in microcentrifuge tube. (2X loading buffer: 100 mM Tris-Cl, pH6.8, 4% sodium dodecyl sulfate, 0.2% bromophenol blue, 20% glycerol, 5% beta-mercapto-ethanol). Cells were lysed in tube and vortexed. Sample was boiled for 10 minutes. 20 microliters of sample was loaded on 7.5%
40 polyacrylamide gel (BioRad) and electrophoresed.


Size fractionated proteins in gel were transferred to nitrocellulose membrane by semidry blotting. Membrane was blocked overnight with 5% powdered milk in phosphate buffer saline (PBS) and 0.05% Tween-20 at 4 degrees C. After 3 washes with PBS/Tween membranes were incubated with

primary antibody for 4 hours at room temperature. After three washes with PBS/Tween membrane was incubated with secondary antibody for 1 hour at room temperature. Finally, a signal was visualized with ECL detection kit from Amersham.

- 5 The compounds of this invention generally show ALK5 receptor modulator activity having IC₅₀ values in the range of 0.0001 to 10 μ M.

Claims:

1. A compound of formula (I) or a pharmaceutically acceptable salt thereof:



(I)

wherein R₁ is naphthyl, anthracenyl, or phenyl optionally substituted with one or more substituents selected from the group consisting of halo, C₁₋₆alkoxy, C₁₋₆alkylthio, C₁₋₆alkyl, -O-(CH₂)_n-Ph, -S-(CH₂)_n-Ph, cyano, phenyl, and CO₂R, wherein R is hydrogen or C₁₋₆alkyl and n is 0, 1, 2 or 3; or R₁ is phenyl fused with an aromatic or non-aromatic cyclic ring of 5-7 members wherein said cyclic ring optionally contains up to two heteroatoms, independently selected from N, O and S;

R₂ is H, NH(CH₂)_n-Ph or NH-C₁₋₆alkyl, wherein n is 0, 1, 2 or 3;

R₃ is CO₂H, CONH₂, CN, NO₂, C₁₋₆alkylthio, -SO₂-C₁₋₆alkyl, C₁₋₆alkoxy, SONH₂, CONHOH, NH₂, CHO, CH₂OH, CH₂NH₂, or CO₂R, wherein R is hydrogen or C₁₋₆alkyl; and

one of X₁ and X₂ is N or CR', and the other is NR' or CHR' wherein R' is hydrogen, OH, C₁₋₆alkyl, or C₃₋₇cycloalkyl; or when one of X₁ and X₂ is N or CR' then the other may be S or O.

2. A compound according to claim 1 wherein R₁ is phenyl optionally substituted with one or more substituents selected from the group consisting of halo, C₁₋₆alkoxy, C₁₋₆alkylthio, and phenyl; or R₁ is phenyl fused with an aromatic or non-aromatic cyclic ring of 5-7 members wherein said cyclic ring optionally contains up to two heteroatoms, independently selected from N, O and S.

3. A compound according to claim 1 or 2 wherein R₂ is positioned ortho to the nitrogen of the pyridyl ring.

4. A compound according to any one of the preceding claims wherein R₃ is CO₂H, CONH₂, CN, NO₂, SONH₂, CONHOH, NH₂, CHO, CH₂OH or CH₂NH₂.

5. A compound according to any one of the preceding claims wherein one of X₁ and X₂ is N or CR', and the other is NR' or CHR' wherein R' is hydrogen, C₁₋₆alkyl, or C₃₋₇cycloalkyl, provided that at least one of X₁ and X₂ is N or NR'; or one of X₁ and X₂ is N and the other is O.

6. A compound according to claim 5 wherein one of X_1 and X_2 is N and the other is NR'.
7. A compound according to any one of the preceding claims wherein each R' is hydrogen.
8. A compound according to claim 1 selected from:
- 4-[4-(4-Fluorophenyl)-5-(2-pyridyl)-1H-imidazol-2-yl]benzonitrile;
 - 4-[4-(4-Fluorophenyl)-5-(2-pyridyl)-1H-imidazol-2-yl]benzonitrile;
 - 4-[4-(4-Fluorophenyl)-5-(2-pyridyl)-1H-imidazol-2-yl]benzoic acid;
 - Methyl 4-[4-(4-fluorophenyl)-5-(2-pyridyl)-1H-imidazol-2-yl]benzoate;
 - Ethyl 4-[4-(4-fluorophenyl)-5-(2-pyridyl)-1H-imidazol-2-yl]benzoate
 - 4-(4-Benzo[1,3]dioxol-5-yl-1-hydroxy-5-pyridin-2-yl-1H-imidazol-2-yl)benzonitrile
 - 4-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1H-imidazol-2-yl)benzonitrile;
 - 4-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1H-imidazol-2-yl)benzoic acid;
 - 2-[4-Benzo[1,3]dioxol-5-yl-2-(4-nitrophenyl)-1H-imidazol-5-yl]pyridine;
 - 3-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1H-imidazol-2-yl)phenylamine;
 - 4-[4-(4-Fluorophenyl)-2-(4-nitrophenyl)-1H-imidazol-5-yl]pyridine;
 - 4-[4-(4-Fluorophenyl)-5-pyridin-2-yl-1H-imidazol-2-yl]phenylamine
 - 4-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1H-imidazol-2-yl)phenyl]methanol;
 - 4-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1H-imidazol-2-yl)benzamide;
 - 4-[4-(2,3-Dihydro-benzo[1,4]dioxin-6-yl)-5-pyridin-2-yl-1H-imidazol-2-yl]benzonitrile;
 - 4-[4-(2,3-Dihydro-benzo[1,4]dioxin-6-yl)-5-pyridin-2-yl-1H-imidazol-2-yl]benzamide;
 - 4-[4-(2,3-Dihydro-benzofuran-5-yl)-5-pyridin-2-yl-1H-imidazol-2-yl]benzamide;
 - 3-[4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1H-imidazol-2-yl]benzonitrile;
 - 4-[4-(2,3-Dihydro-benzofuran-6-yl)-5-pyridin-2-yl-1H-imidazol-2-yl]benzonitrile;
 - 4-[4-(2,3-Dihydro-benzofuran-6-yl)-5-pyridin-2-yl-1H-imidazol-2-yl]benzamide;
 - 3-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1H-imidazol-2-yl)benzoic acid;
 - 4-[4-(4-Methoxyphenyl)-5-(2-pyridyl)-1H-imidazol-2-yl]benzonitrile;
 - 4-[4-(2,2-Difluoro-benzo[1,3]dioxol-5-yl)-5-pyridin-2-yl-1H-imidazol-2-yl]benzamide;
 - 4-[4-(2,3-Dihydro-benzo[1,4]dioxin-6-yl)-1-methyl-5-pyridin-2-yl-1H-imidazol-2-yl]benzamide;
 - 4-[5-(2,3-Dihydro-benzo[1,4]dioxin-6-yl)-1-methyl-4-pyridin-2-yl-1H-imidazol-2-yl]benzamide;
 - 4-(5-Benzo[1,3]dioxol-5-yl-4-pyridin-2-yl-oxazol-2-yl)benzonitrile;
 - 4-(5-Benzo[1,3]dioxol-5-yl-4-pyridin-2-yl-oxazol-2-yl)benzamide; and
 - 4-(4-Benzo[1,3]dioxol-5-yl-5-pyridin-2-yl-1H-pyrrol-2-yl)benzamide;
- or a pharmaceutically acceptable salt thereof.

9. A pharmaceutical composition comprising a compound according to any one of the claims 1 to 8, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or diluent.
10. A method of inhibiting the TGF- β signaling pathway in mammals, comprising administering to a mammal in need of such treatment, a therapeutically effective amount of a compound according to any one of claims 1 to 8, or a pharmaceutically acceptable salt thereof.
11. A method for treating a disease selected from chronic renal disease, acute renal disease, wound healing, arthritis, osteoporosis, kidney disease, congestive heart failure, ulcers, ocular disorders, corneal wounds, diabetic nephropathy, impaired neurological function, Alzheimer's disease, trophic conditions, atherosclerosis, peritoneal and sub-dermal adhesion, any disease wherein fibrosis is a major component, and restenosis, comprising administering to a mammal in need of such treatment, a therapeutically effective amount of a compound according to any one of claims 1 to 8, or a pharmaceutically acceptable salt thereof.
12. A method for inhibiting matrix formation in mammals, comprising administering to a mammal, a therapeutically effective amount of a compound according to any one of claims 1 to 8, or a pharmaceutically acceptable salt thereof.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/09147

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/04 C07D413/04 C07D405/14 A61K31/4178 A61P19/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 656 644 A (J. LEROY ADAMS) 12 August 1997 (1997-08-12) cited in the application column 12 -column 15; claims; examples 1-97	1,2,6,7, 9
A	US 5 776 954 A (S.E. DE LASZLO ET AL.) 7 July 1998 (1998-07-07) the whole document	1,2,9
A	GB 1 381 031 A (CIBA-GEIGY) 22 January 1975 (1975-01-22) page 7; claims	1,2,6,7, 9
A	WO 95 13067 A (SMITHKLINE-BEECHAM) 18 May 1995 (1995-05-18) claims 1,9; examples 5,11,12	1,2,9
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

31 August 2000

Date of mailing of the international search report

15/09/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Francois, J

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/09147

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 22109 A (MERCK) 28 May 1998 (1998-05-28) the whole document	1,2,6,7, 9
P,A	CHEMICAL ABSTRACTS, vol. 131, no. 7, 1999 Columbus, Ohio, US; abstract no. 87659s, page 641; XP002146305 abstract & JP 11 180958 A (YAMANOUCHI) 6 July 1999 (1999-07-06)	1,2,6,7, 9

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/09147

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5656644	A	12-08-1997	US 5916891 A	29-06-1999
US 5776954	A	07-07-1998	NONE	
GB 1381031	A	22-01-1975	CH 579072 A	31-08-1976
			CH 561202 A	30-04-1975
			AR 194594 A	31-07-1973
			AR 198820 A	24-07-1974
			AR 198658 A	15-07-1974
			AR 195910 A	15-11-1973
			AT 319939 B	10-01-1975
			AT 319940 B	10-01-1975
			AT 319941 B	10-01-1975
			AT 316544 B	15-06-1974
			AU 472065 B	13-05-1976
			AU 4205572 A	15-11-1973
			BE 783244 A	10-11-1972
			CA 1012148 A	14-06-1977
			CH 561716 A	15-05-1975
			CH 561717 A	15-05-1975
			CH 561718 A	15-05-1975
			CS 181711 B	31-03-1978
			CS 181745 B	31-03-1978
			CS 181746 B	31-03-1978
			CS 181747 B	31-03-1978
			CY 936 A	23-06-1978
			DD 97654 A	12-05-1973
			DE 2221546 A	16-11-1972
			ES 402563 A	01-11-1975
			FI 57407 B	30-04-1980
			FR 2137740 A	29-12-1972
			HK 61477 A	16-12-1977
			HU 164884 B	28-05-1974
			IE 36319 B	13-10-1976
			IL 39285 A	10-02-1975
			JP 56030352 B	14-07-1981
			KE 2796 A	06-01-1978
			MY 11878 A	31-12-1978
			NL 7206346 A	14-11-1972
			NO 137445 B	21-11-1977
			SE 405731 B	27-12-1978
			SU 456409 A	05-01-1975
			SU 502605 A	05-02-1976
			SU 489329 A	25-10-1975
			SU 489330 A	25-10-1975
			US 3940486 A	24-02-1976
			US 3929807 A	30-12-1975
			YU 17680 A	30-04-1981
			YU 83779 A, B	31-12-1980
			YU 123372 A, B	31-12-1980
			YU 264478 A, B	31-12-1980
			YU 264578 A, B	31-12-1980
			YU 264678 A, B	31-12-1980
WO 9513067	A	18-05-1995	EP 0727998 A	28-08-1996
			JP 9505055 T	20-05-1997
WO 9822109	A	28-05-1998	AU 5442898 A	10-06-1998

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/09147

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9822109 A		EP 0959886 A	01-12-1999
		US 5880139 A	09-03-1999
JP 11180958 A	06-07-1999	NONE	